

On the origin of Fe/S cluster biosynthesis in eukaryotes

Anastasios D. Tsaousis^{1*}

¹School of Biosciences, University of Kent, United Kingdom

Submitted to Journal:
Frontiers in Microbiology

Specialty Section:
Evolutionary and Genomic Microbiology

Article type:
Hypothesis and Theory Article

Manuscript ID:
482978

Received on:
04 Jul 2019

Revised on:
27 Sep 2019

Frontiers website link:
www.frontiersin.org

Conflict of interest statement

The authors declare that the research was conducted in the absence of any commercial or financial relationships that could be construed as a potential conflict of interest

Author contribution statement

ADT has conceptualised and wrote the manuscript

Keywords

LECA (Last Eukaryotic Common Ancestor), iron sulfur cluster biogenesis, SUF machinery, Isc (iron sulfur cluster) machinery, eukaryotic evolution, CIA machinery

Abstract

Word count: 188

Iron and sulfur are indispensable elements of every living cell, but on their own these elements are toxic and require dedicated machineries for the formation of Fe/S clusters. In eukaryotes, proteins requiring Fe/S clusters (Fe/S proteins) are found in or associated with various organelles including the mitochondrion, endoplasmic reticulum, cytosol and the nucleus. These proteins are involved in several pathways indispensable for the viability of each living cell including DNA maintenance, protein translation and metabolic pathways. Thus, the formation of Fe/S clusters and their delivery to these proteins has a fundamental role in the functions and the evolution of the eukaryotic cell. Currently, most eukaryotes harbor two (located in cytosol and mitochondrion) or three (located in plastid) machineries for the assembly of Fe/S clusters, but certain anaerobic microbial eukaryotes contain Sulfur Mobilization (SUF) machineries that were previously thought to be present only in archaeal lineages. These machineries could not only stipulate which pathway was present in the last eukaryotic common ancestor (LECA), but they could also provide clues regarding presence of an Fe/S cluster machinery in the proto-eukaryote and evolution of Fe/S cluster assembly machineries in all eukaryotes.

Contribution to the field

Iron and sulphur are indispensable elements of every living cell, but on their own are toxic and require dedicate and indispensable machineries for the formation of Fe/S clusters. In eukaryotes, proteins requiring Fe/S clusters (Fe/S proteins) are found in or associated with various organelles including the mitochondrion, endoplasmic reticulum, cytosol and the nucleus. These proteins are involved in several pathways indispensable for the viability of each living cell including metabolic pathways, DNA maintenance and protein translation. Thus, the formation and delivery of the Fe/S clusters to these proteins has fundamental role in the functions and the evolution of the eukaryotic cell. Over the last decade there have been significant discoveries in regards to the evolution of eukaryotes and the role of the Fe-S biosynthetic pathways in their adaptations to unique lifestyles. Currently, most eukaryotes harbour two (located in cytosol and mitochondrion) or three (located in plastid) machineries for the assembly of Fe/S clusters. We will present a small summary of these machineries and their roles within the eukaryotic cell. Despite this, certain anaerobic microbial eukaryotes contain machineries that were previously thought to be commonly found in archaeal lineages. Which these machineries are and how have they been acquired or preserved in these various eukaryotic lineages? We will present these exemptions and then we will focus on the Sulphur Mobilization (SUF) machinery, which is commonly found in plastids, but also in the cytosol and/or mitochondria in various anaerobic/microaerophilic protists such as *Blastocystis*, *Pygmaia* and *Stygiella*. This machinery is considered to be the most “ancient” Fe-S cluster machinery (not only in eukaryotes). We will provide alternative theories/scenarios based on current published data regarding the presence, function and evolution of this machinery and co-evolution with other machineries in eukaryotes. The presence of the SUF machinery in various eukaryotes could not only stipulate which pathway could have been present in the last eukaryotic common ancestor, but they could also provide clues into the evolution of Fe/S cluster assembly machineries in eukaryotes. Based on current data, we will propose various scenarios on the evolution of the Fe-S cluster machineries in eukaryotes and we will suggest that a SUF-like ancient Fe/S cluster machinery could have been present in proto-eukaryotic cell or the last common eukaryotic ancestor. This is timely, due to the various recent publications on sequencing the genomes of various lineages of Asgard archaea in an attempt to identify the nature of the “founding lineage” of eukaryotes. Based on the proposed scenarios that will be discussed in this article, such a lineage, will provide us with insights on the presence and function of a fundamental biosynthetic pathway such as the Fe-S cluster biosynthesis. Such an essential pathway has yet to be discovered in these archaeal lineages; but according to Prof. Thijs Ettema (presentation in a recent conference) many more archaeal lineages are soon to be published, and thus will open a new field of explorations, while providing a hypothesis to be tested.

Ethics statements

Studies involving animal subjects

Generated Statement: No animal studies are presented in this manuscript.

Studies involving human subjects

Generated Statement: No human studies are presented in this manuscript.

Inclusion of identifiable human data

Generated Statement: No potentially identifiable human images or data is presented in this study.

In review

Data availability statement

Generated Statement: No datasets were generated or analyzed for this study.

In review

Hypothesis & Theory:

On the origin of Fe/S cluster biosynthesis in eukaryotes

Anastasios D. Tsaousis¹

1. Laboratory of Molecular and Evolutionary Parasitology, RAPID group, School of
Biosciences, University of Kent, Canterbury, Kent, CT2 7NJ, United Kingdom

*corresponding author: Dr. Anastasios Tsaousis (A.Tsaousis@kent.ac.uk)

Abstract

Iron and sulfur are indispensable elements of every living cell, but on their own these elements are toxic and require dedicated machineries for the formation of Fe/S clusters. In eukaryotes, proteins requiring Fe/S clusters (Fe/S proteins) are found in or associated with various organelles including the mitochondrion, endoplasmic reticulum, cytosol and the nucleus. These proteins are involved in several pathways indispensable for the viability of each living cell including DNA maintenance, protein translation and metabolic pathways. Thus, the formation of Fe/S clusters and their delivery to these proteins has a fundamental role in the functions and the evolution of the eukaryotic cell. Currently, most eukaryotes harbor two (located in cytosol and mitochondrion) or three (located in plastid) machineries for the assembly of Fe/S clusters, but certain anaerobic microbial eukaryotes contain Sulfur Mobilization (SUF) machineries that were previously thought to be present only in archaeal lineages. These machineries could not only stipulate which pathway was present in the last eukaryotic common ancestor (LECA), but they could also provide clues regarding presence of an Fe/S cluster machinery in the proto-eukaryote and evolution of Fe/S cluster assembly machineries in all eukaryotes.

Introduction

Iron/sulfur (Fe/S) clusters are fundamental and ubiquitous factors. All living cells have biosynthetic machineries responsible for their assembly and delivery, since the individual components (iron and sulfur; Fe and S) are toxic for the cells themselves (Lill et al. 1999; Lill 2009). Importantly, Fe/S clusters are essential factors of proteins involved in essential functions of the cell including, but not restricted to, photosynthesis, respiration, DNA replication and repair, and regulation of gene expression (Lill et al. 2012). Eukaryotes are not the exception to this paradigm. The typical Fe/S biosynthetic machineries found in bacteria and archaea have also been identified in eukaryotes, but compartmentalization and evolution of these machineries in several eukaryotes are still under investigation. A typical eukaryotic cell harbors the Iron-Sulfur Cluster (ISC) in the mitochondria and the Cytosolic Iron/Sulfur cluster Assembly (CIA) machinery in the cytosol, while plastid-carrying cells also harbor the Sulfur Mobilization (SUF) machinery in their plastids.

Among those, the ISC machinery has been considered to be the reason for the existence of mitochondria (Hjort et al. 2010; Lill et al. 1999; Lill 2009), and fundamental for the evolution of eukaryotes. Nonetheless, what happens when a eukaryote does not harbor any mitochondria (Karnkowska et al. 2016)? Could this organism provide some clues about the presence of Fe/S biosynthetic machineries in the early eukaryotes and their role in the evolution of the eukaryotic cell?

Fe-S cluster assembly in mitochondrial diversity

It is widely accepted that mitochondria originated from or within the alpha-proteobacteria (Gawryluk 2018; Gray et al. 1999; Gray et al. 2001; Martijn et al. 2018), whereby the latter was “engulfed” by a eukaryotic host and potentially gave rise to the Last Eukaryotic Common Ancestor (LECA). Nevertheless, questions regarding why, how and when this event took place are still under debate (Embley and Martin 2006; Gabaldon 2018; Gray et al. 2001; Lane and Martin 2015; Lane and Martin 2016; Martin et al. 2016; Pittis and Gabaldon 2016). It is quite apparent from accumulated data that the acquisition of mitochondria has been the decisive step in eukaryogenesis (Martin et al. 2016). One hypothesis postulates that the mitochondria fulfilled energy requirements of the cell thus their

presence provided a selective advantage to the organisms bearing them to become eukaryotes (Pittis and Gabaldon 2016; Lane and Martin 2015; Lane and Martin 2016). Another hypothesis, which does not exclude others, suggests that the reason for the existence of mitochondria could have been the assembly of Fe/S clusters (Lill et al. 1999), the latter being the only mitochondrial biosynthetic pathway that is essential for survival of eukaryotic cells. So far, this has been shown experimentally in yeast (Braymer and Lill 2017), mammalian cells (Rouault and Maio 2017) and trypanosomes (Pena-Diaz and Lukes 2018).

Further support to this hypothesis arose from investigations in previously considered “primitive” amitochondriate eukaryotes. These organisms were shown to harbor mitochondrial-related organelles (MROs), a secondarily reduced form of mitochondria, including hydrogen producing organelles called hydrogenosomes in *Trichomonas* (Muller 1973), or highly reduced remnant organelles called mitosomes, which were found in *Giardia* (Tovar et al. 2003); microsporidia (Tsaousis et al. 2008; Williams et al. 2002) and *Entamoeba* (Tovar et al. 1999). Whether a “primitive” amitochondriate eukaryote could exist or not, is still under debate (Margulis et al. 2006). Nonetheless, a eukaryote that secondarily lost its mitochondria was identified recently (Karnkowska et al. 2016). Interestingly, the only biosynthetic pathway conserved in all these organelles is the assembly of Fe/S clusters, providing further support on the necessity/importance of this machinery for cell viability. From an evolutionary standpoint, it will be important to elucidate how the eukaryotic cell supported its needs for Fe/S clusters, before the acquisition of mitochondria. To provide insight on this matter, I will first need to examine the distribution of various Fe/S cluster machineries in eukaryotic cells and their necessity to the host’s functions, followed by various theories on the evolution of Fe-S cluster machineries across eukaryotes.

Mitochondrial Fe/S cluster machinery

All mitochondria investigated so far possess some semblance of an Fe/S cluster biosynthetic pathway for *de novo* assembly of Fe/S clusters into organellar apo-proteins (see below), but potentially for the support of cytosolic and nuclear apo-proteins as well (Ali and Nozaki 2013; Lill 2009). The typical mitochondrial machinery is the Iron-Sulfur Cluster (ISC), which is comprised of 18 (currently known in yeast) proteins (Braymer and Lill 2017), all of which are involved in the biogenesis and

trafficking of clusters in mitochondria (**Figure 1**). The process is divided into four stages (for detailed review see Braymer and Lill, 2017): (i) *de novo* [2Fe-2S] cluster synthesis; (ii) trafficking of [2Fe-2S] clusters and insertion into mitochondrial apo-proteins, or mitochondrial export of an as yet unknown Sulfur-containing species (X-S) to the cytosol; (iii) conversion of [2Fe-2S] into [4Fe-4S] clusters; and lastly (iv) trafficking of [4Fe-4S] clusters and insertion into mitochondrial [4Fe-4S] apo-proteins (e.g. lipoate synthase, succinate dehydrogenase, components of respiratory complex I). Most organisms harboring mitochondria encode some of these components, including organisms with remnant mitochondria such as *Giardia* (Tovar et al. 2003), *Cryptosporidium* (Miller et al. 2018) and microsporidia (Freibert et al. 2017; Goldberg et al. 2008), in which ISC stages iii and iv are lacking ([4Fe-4S] cluster synthesis & targeting; **Figure 1**), due to the lack of mitochondrial apo-proteins requiring [4Fe-4S] clusters.

Cytosolic Fe/S cluster machinery

All eukaryotes require a cytosolic Fe/S cluster (CIA) machinery to support cytosolic and nuclear Fe/S cluster proteins (Tsaousis et al. 2014). So far, 11 proteins have been identified in both mammals and yeast as responsible for synthesis, trafficking and insertion of clusters in the cytosol and the nucleus (Braymer and Lill 2017; Tonini et al. 2018). Of these, several CIA protein complexes support different stages in the process (**Figure 2a**). For example, a bridging [4Fe-4S] cluster is assembled on the Cfd1-Nbp35 complex, which depends on the as yet unidentified molecule X-S from the mitochondrial ISC machineries. Subsequently, the electron transfer chain from NADPH via the diflavin reductase Tah18 and the Fe/S protein Dre2 is required. In the next phase, the transiently bound [4Fe-4S] cluster of Cfd1-Nbp35 is transferred to and inserted into apo-proteins by the Fe/S protein Nar1, and the CIA targeting complex consisting of Cia1, Cia2 and Mms19 (Stehling et al. 2012; Stehling et al. 2013). This entity also binds the Lto1-Yae1 adapter complex via a conserved C-terminal tryptophan in Lto1 to recruit the ABC protein Rli1 (participates in ribosome assembly and ribosome recycling) for dedicated assembly of its two [4Fe-4S] clusters (Lill et al. 2015; Paul et al. 2015). The CIA machinery may also support ATP-dependent DNA helicases such as Rad3, XPD, FANCI, and RTEL1, which are involved in DNA damage repair and telomere maintenance (Rudolf et al. 2006). Interestingly, mitochondria or related organelles,

such as hydrogenosomes and mitosomes (see above) seem to be essential for the support of the CIA machinery in the biogenesis of cytosolic and nuclear Fe/S clusters (Freibert et al. 2017; Stehling et al. 2014; Tsaousis et al. 2014). Despite this, organisms harboring these “reduced” mitochondria appear to lack certain components of the CIA machinery (e.g. Tah18, Dre2 and Cfd1) that are otherwise essential in mammals and yeast (Tsaousis et al. 2014; Vacek et al. 2018). Even more intriguingly, microbial organisms such as cryptophytes and chlorarachniophytes that harbor cytosols from two organisms (main and cytosol of their phototrophic symbiont), seem to have two diverse and functional CIA machineries – one in each compartment – which are supported by their corresponding organelles (Grosche et al. 2018).

Plastid Fe/S cluster machinery

Apo-proteins in plastids and plastid-related organelles are supported by the Sulfur mobilization (SUF) machinery, which was acquired from Cyanobacteria. The six major proteins that encompass the bacterial-type SUF machinery are also present in plastids (SufA, SufB, SufC, SufD, SufE and SufS; **Figure 3a**), one of which (SufC) is commonly encoded by the plastid genome (Le Corguille et al. 2009). Using genetic and biochemical investigations in prokaryotes it was shown that SufE and SufS are involved in the Sulfur mobilization from cysteine, while SufB, SufC and SufD form a complex where SufB harbors both the *de novo* assembled Fe/S clusters and a flavin redox cofactor (Couturier et al. 2013). However, recent experimental structural studies have shown a dynamic motion of the SufB₁-SufC₂-SufD₁ complex, that could be universally applicable to all the SUF systems, including the archaeal SufB₂-SufC₂ complex (Hirabayashi et al. 2015) (discussed below). In addition, SufA could act as a carrier protein, along with numerous other carrier proteins that are currently found [(Fontecave et al. 2005; Wollers et al. 2010), for review see Couturier et al., 2013]. As such, the plastidial Fe/S assembly machinery has been mostly characterised in *Arabidopsis thaliana*, where 15 proteins have been experimentally localized and one of which (SufSE) was shown to be targeted in both the plastids and mitochondria (Balk and Pilon 2011; Couturier et al. 2013). To that end, the plastidial Fe/S cluster assembly is responsible for the support of housekeeping apo-proteins of the organelle and currently is unclear if it can support the CIA machinery in cytosol of the cells (similar to the ISC machinery).

Fe/S cluster assembly in amitochondriates

The discovery of a eukaryote that secondarily lost its mitochondria (Karnkowska et al. 2016), raises the question of Fe/S cluster biosynthesis in this organism, since this is the only biosynthetic function found in all mitochondria-related organelles investigated so far (Hjort et al. 2010; Santos et al. 2018). The oxymonad *Monocercomonoides* sp. [currently named *M. exilis* (Treitli et al. 2018)] is the first eukaryotic organism with no microscopic evidence for the existence of a mitochondrion. This finding was further supported by extensive genome surveys that failed to find any mitochondrial proteins, including homologues of the mitochondrial ISC pathway (Karnkowska et al. 2016). Despite this, the genome of *Monocercomonoides* does encode components of the CIA machinery (**Figure 2c**), in addition to homologues of a SUF system (**Figure 3a,b**). The origin of these SUF homologues though unclear, seems to be bacterial (Karnkowska et al. 2016) (see below). Due to the lack of an *in situ* transfection system, *Monocercomonoides* SufC and SufB homologues were heterologously expressed in *Trichomonas vaginalis* and *Saccharomyces cerevisiae*, whereby they both localized in the cytosol of both organisms (Karnkowska et al. 2016).

Recent investigations by Vacek et al (2018) demonstrated that oxymonads and organisms (Preaxostyla group, Metamonada, Excavata) related to *M. exilis* also harbor a SUF machinery (Vacek et al. 2018). Genomic and transcriptomic surveys have shown the presence of components of the SUF machinery in six additional closely related species, suggesting that transition from ISC to SUF preceded the last common ancestor of the lineage (Vacek et al. 2018). A follow-up inventory of all the homologues of the CIA machinery in these organisms showed that its major components are still present, consistent with previous observations that the lack of mitochondria or more specifically of the ISC machinery did not have any effect in the maturation of cytosolic Fe/S proteins (Vacek et al. 2018).

Exceptions to the *status quo* (alternative directions)

1. The case of *Entamoeba* and *Mastigamoeba*

In addition to the machineries described above, some organisms have acquired new processes for the *de novo* assembly of their Fe/S clusters. The genomes of the amoebozoans *Entamoeba histolytica* and *Mastigamoeba balamuthi* (both thriving in low-oxygen environments) do not encode any components of

the ISC machinery and instead they harbor a Nitrogen Fixation (NIF) machinery that was laterally acquired from an epsilon proteobacterion (Ali et al. 2004; van der Giezen et al. 2004). Components of the machinery were shown to localize in the mitosome of *E. histolytica* (Maralikova et al. 2010) [though this is still under debate (Nyvltova et al. 2013)], while replica components of *M. balamuthi* were shown to localize in both the cytosol and its hydrogenosomal-like structures (Nyvltova et al. 2013). It is still unclear whether the function of a NIF system could be more advantageous over the ISC system, but it seems to be the “preferred” way in this lineage. Despite this alteration, components of the CIA machinery are present in both organisms (Pyrih et al. 2016; Tsaousis et al. 2014) (with the exception of Tah18, Dre2 and Cfd1), suggesting that ISC machinery might not [as previously thought (Lill et al. 1999)] be indispensable for the function of the CIA machinery.

2. The case of *Blastocystis*, *Pygusua*, *Stygiella* and others?

Blastocystis is an obligatory anaerobic stramenopile. *Blastocystis* was the first non-photosynthetic eukaryotic organism to be shown to encode an ancient SUF system (Tsaousis et al. 2012), in addition to an ISC machinery that is localized in mitochondria (Tsaousis et al. 2012) and a CIA machinery that is localized in the cytosol (Tsaousis et al. 2014). The SUF system of *Blastocystis* is similar to the one of Methanomicrobiales in that both display fusion of the *SufC* and *SufB* genes. Phylogenetic analysis showed that both *Blastocystis* homologues grouped with those of the archaea into a strongly supported clade, indicating lateral acquisition of the gene from Methanomicrobiales (Tsaousis et al. 2012). The fused gene is found in the genomes of all *Blastocystis* subtypes, in addition to the genome of *Proteromonas lacertae* (found in BioProject: PRJNA386230), a Stramenopile species closely related to *Blastocystis*. Functional characterization of the *Blastocystis* protein showed that it binds [4Fe-4S] clusters and has ATPase activity. The protein was shown to localize in the cytosol of the parasite and to be overexpressed under oxygen-stressed conditions (Tsaousis et al. 2012). This was unsurprising, since in various bacteria, it has been demonstrated that the machinery is overexpressed under oxygen stress or iron depletion conditions, in order to support the potentially damaged apo-proteins of the cell (Metttert et al. 2008; Rangachari et al. 2002).

Following its discovery in *Blastocystis*, a fused *SufCB* gene was later found in other distantly related microbial eukaryotes. The first was the breviate *Pygсуia biforma*, a free-living anaerobe, but aerotolerant amoeboid flagellate isolated from hypoxic marine sediments. The organism branches at the base of the eukaryotic supergroup Obazoa, which is comprised of animals, fungi and apusomonads (**Figure 3b**). The *P. biforma* genome encodes two homologues of the protein (Stairs et al. 2014). Localization experiments showed that one homologue localizes in mitochondria, while the other localizes in the cytosol (Stairs et al. 2014). Phylogenetic analysis showed that both *P. biforma* homologues branch closely with those of *Blastocystis*. Interestingly, analysis of the RNA-seq data did not show expression of any of the components of the mitochondrial ISC machinery, while components of the CIA machinery (Cia1, Nbp35, Cfd1, Nar1, Cia2, and Met18) were present (Stairs et al. 2014).

A fused *SufCB* gene was also found in *Stygiella incarcerata* along with genes encoding components of the mitochondrial ISC machinery (Leger et al. 2016). *Stygiella incarcerata* a microaerophilic jakobid flagellate inhabiting anoxic environments and is distantly related to Stramenopiles and Breviata (e.g. *Blastocystis* and *Pygсуia* respectively; **Figure 3b**). The *SUF*CB gene of *S. incarcerata* displayed the same characteristics as the homologues of *Blastocystis* and *Pygсуia*, and it lacked mitochondrial targeting peptides suggesting a potential cytosolic localization. While the authors did not find any introns in the transcriptome derived fused gene, data from the closely related jakobid *Velundella trypanoides* (found in BioProject: PRJNA268717) also demonstrated the presence of a homologue (Leger et al. 2016), suggesting that the gene is likely not a contaminant. Phylogenetic analysis showed that the *SUF* eukaryotic homologues from *Blastocystis*, *Pygсуia* and *Stygiella* formed a strongly supported clade, with Methanomicrobiales as a well-supported sister group (Leger et al. 2016), consistent with previous observations (Stairs et al. 2014; Tsaousis et al. 2012). How is it possible for organisms that are so distantly related to have a *SUF*CB homologue?

Various scenarios could explain the presence of this machinery in at least three eukaryotic lineages. Herein, I will discuss three scenarios (**Figure 3c-g**) while providing pros and cons for each hypothesis:

1st Theory:

All three organisms (or their ancestors) acquired the methanoarchaeal SufCB independently, likely while inhabiting the same environmental niche (**Figure 3c**). This scenario suggests three independent transfers: once in the common ancestor of *Blastocystis* and *Proteromonas*, once in *Stygiella* and once in *Pygsuia*. Each transfer would require co-existence of the donor lineage with each eukaryote separately. Consequently, this setting implies that the ancestors of these organisms co-habitated in similar environments with Methanomicrobiales, which allowed for transfer and incorporation of genes in their genomes. The intriguing question, under this scenario, is why only a single fused gene was transferred or incorporated from these methanomicrobes in the genomes of diverse protozoa lineages (Tsaousis et al. 2012)?

2nd Theory:

The methanoarchaeal SufCB gene was acquired by one of the three eukaryotic organisms (or their ancestors) and then laterally transferred to the others (**Figure 3d, e & f**). It is well established that lateral gene transfer events from eukaryotes to eukaryotes are not as uncommon as it was once thought (Danchin 2016; Eme et al. 2017; Leger et al. 2018). This type of scenario requires that at least two of the protists co-habited with the donor lineage in the same or similar niches at some point of their life cycles. For example, *Blastocystis* and *Proteromonas* spend the majority of their life cycle in the gut of various organisms. Nonetheless, *Blastocystis* is excreted in the environment as a cyst. If cysts were shed in hypoxic environments, then the possibility of *Pygsuia* and *Stygiella* encountering *Blastocystis* (or its ancestor) and subsequently exchanging genetic material is not entirely far-fetched. Interestingly, with the exception of the SufCB gene, to our knowledge, no other genes share the same origins (or clustering) in these three groups.

3rd Theory:

The methanoarchaeal SufCB was present in the last eukaryotic common ancestor (LECA) (**Figure 3g**). The LECA had to have a machinery for the assembly of Fe-S clusters to support its apo-proteins, even before the acquisition of the alpha-proteobacterium that gave rise to the present-day mitochondria. Notably, it has been suggested that the CIA machinery, which is present in all eukaryotes

investigated so far is a eukaryotic innovation (Freibert et al. 2017; Tsaousis et al. 2014). Since the ISC machinery is found only in mitochondria and the NIF machinery is only present in two closely related organisms, it is unlikely that either one was present in LECA. Thus, an ancestral SUF machinery, which is commonly found in archaea (Outten 2015), could have been present in LECA. Considering that *SufCB* is not only the most “ancient machinery” (Tokumoto et al. 2004) amongst all biosynthetic apparatuses, but also the most widespread across lineages, it is plausible that the SufCB was present in the common ancestor of eukaryotes as well. The machinery could have either been acquired by a methanoarchaeon or it could have been present in the archaeal group that gave rise to modern eukaryotes (Eme et al. 2018; Spang and Ettema 2017; Zaremba-Niedzwiedzka et al. 2017). This scenario could explain the presence of a biosynthetic machinery in three distantly related eukaryotic lineages, but it also infers multiple losses of this machinery in the rest of the lineages. Under this scenario, the case of oxymonads is of interest (Karnkowska et al. 2016; Vacek et al. 2018). How can a separate origin of SUF be explained? One explanation would be that the ancestrally acquired SUF was lost and a SUF of different origin was acquired upon loss of mitochondria. Thus, I hypothesize that eukaryotes maintain the chassis that would allow reacquisition of SUF-like machinery. This hypothesis could be tested by incorporating the eukaryotic SUF machineries in various model organisms across the eukaryotic tree of life (e.g. *Saccharomyces*, *Trypanosoma*, *Tetrahymena*, *Dictyostelium*). It’s worth mentioning that the 3rd theory does not necessary exclude the other theories above.

Discussion: Fe/S cluster biosynthesis during the evolutionary history of eukaryotes

Given the discovery of this fused gene in diverse lineages of eukaryotes, speculative scenarios propose an initial transfer of the SufCB from an archaeal source into an ancestral microbial eukaryote (**Figure 3c,g**), and/or lateral gene transfer events to other eukaryotes (Leger et al. 2016; Tsaousis et al. 2014) (**Figure 3d-f**). Nevertheless, it is imperative to highlight the importance of this pathway in the evolution and adaptation of eukaryotes.

The last eukaryotic common ancestor (LECA) lived about 1.8 billion years ago (Betts et al. 2018) and seems to have been more complicated than was previously thought (Koonin 2015). It has been speculated that LECA contained organelles and functions that even mirror some of the current microbial

eukaryotes, based on comparative genomic analyses with the closest archaeal-relative lineage, the Lokiarchaeota (Eme and Ettema 2018; Eme et al. 2018; Spang et al. 2015; Spang et al. 2017; Spang et al. 2018; Zaremba-Niedzwiedzka et al. 2017). Among those, it is currently suggested that LECA possessed mitochondria, endomembrane system along with nucleus, actin cytoskeleton, endocytosis and/or phagocytosis and a ubiquitin network (Akil and Robinson 2018; Embley and Williams 2015; Eme and Ettema 2018; Koonin 2015; Spang et al. 2015). Metabolically, based on investigations in Lokiarchaeota, LECA could have been transitioning from anaerobic to aerobic metabolism (due to the acquisition of the mitochondria; aerobic respiration) with a potentially hydrogen-dependent autotrophic lifestyle (Martin et al. 2016; Sousa et al. 2016). Some of these pathways need enzymes (apo-proteins) that require Fe/S clusters in order to function, including DNA/RNA polymerases and anaerobic proteins (e.g. pyruvate ferredoxin oxidoreductase; PFO), which have been identified in Lokiarchaeota (Sousa et al. 2016). LECA must have harbored a biosynthetic pathway to support the assembly and trafficking of these Fe/S clusters. The presence of a SUF-like machinery in LECA is plausible, since it is the most common machinery amongst archaeal lineages and is also not compartmentalized in most eukaryotes (Karnkowska et al. 2016; Leger et al. 2016; Stairs et al. 2014; Tsaousis et al. 2012). Footprints of this ancient machinery still remain in modern eukaryotes and it is not an invalid prediction that more organisms having this machinery will be discovered. Whether the machineries that are present in *Blastocystis/Proteromonas*, *Pygmaia* and *Stygiella* lineages are remnants of the initial machinery (LECA) or later acquisitions (see scenarios **Figure 3c-g**) will need further investigations; current data clearly illustrate that the CIA and SUF-like machineries can clearly co-exist (Karnkowska et al. 2016; Leger et al. 2016; Stairs et al. 2014; Tsaousis et al. 2012; Tsaousis et al. 2014; Vacek et al. 2018).

It is also important to note that SUF-like machineries have been shown to be upregulated under oxygen stress conditions to support the potential degradation of Fe-S clusters of proteins (Metttert et al. 2008; Rangachari et al. 2002). This function/support would have been essential during the transformation of proto-eukaryotic cells to LECA, since during that period there would have been a transition to increasing concentrations of oxygen (Lane and Martin 2016). A SUF-like machinery would have been able to compensate for the potential damage of Fe/S clusters from oxygen allowing cells to slowly adjust to their new environments. In parallel, acquisition of mitochondria provided not only an oxygen

protective compartment for the formation of Fe/S clusters, but also the ISC machinery as well (Lill et al. 1999; Lill et al. 2015). Later on, adaptation of these cells to oxygen rich environments and expansion of the CIA machinery in the cytosol along with its ability to “communicate” with the mitochondrial ISC machinery (e.g. ATM1 for transfer of X-factor; **Figures 1 & 2**), resulted into the SUF-like machinery becoming redundant to the ancestors of most eukaryotic lineages. Eukaryotes that still remained under oxygen depleted conditions either retained the SUF-like machinery (scenario **Figure 3g**) or later acquired a homologue of this (Vacek et al. 2018).

Here, I propose various scenarios on the evolution of the Fe-S cluster machineries in eukaryotes and I suggest that a SUF-like ancient Fe/S cluster machinery could have been present in the proto-eukaryotic cell or LECA. Current ‘omics data do not provide an answer to this question, but existing efforts to broadly sample the large diversity of archaeal and eukaryotic lineages could provide the missing pieces of this unsolved puzzle.

Acknowledgments

ADT was supported by an internal grant from the University of Kent and BBSRC research grant (BB/M009971/1). I would like to thank Dr. Eleni Gentekaki for her constructive comments and critique on the manuscript. I would like to thank Dr. Joel Dacks for providing me the phylogenetic tree showing the relationship between eukaryotes that was used in Figure 3.

References:

- Akil C. and Robinson R. C. (2018). Genomes of Asgard archaea encode profilins that regulate actin. *Nature* 562, 439-443. doi: 10.1038/s41586-018-0548-6 [doi].
- Ali V. and Nozaki T. (2013). Iron-sulfur clusters, their biosynthesis, and biological functions in protozoan parasites. *Advances in Parasitology* 83, 1-92. doi: 10.1016/B978-0-12-407705-8.00001-X [doi].

353 Ali V., Shigeta Y., Tokumoto U., Takahashi Y. and Nozaki T. (2004). An intestinal parasitic protist,
 354 *Entamoeba histolytica*, possesses a non-redundant nitrogen fixation-like system for iron-sulfur cluster
 355 assembly under anaerobic conditions. *The Journal of biological chemistry* 279, 16863-16874. doi:
 356 10.1074/jbc.M313314200 [doi].

357 Balk J. and Pilon M. (2011). Ancient and essential: the assembly of iron-sulfur clusters in plants. *Trends*
 358 *in plant science* 16, 218-226. doi: 10.1016/j.tplants.2010.12.006 [doi].

359 Betts H. C., Puttick M. N., Clark J. W., Williams T. A., Donoghue P. C. J. and Pisani D. (2018).
 360 Integrated genomic and fossil evidence illuminates life's early evolution and eukaryote origin. *Nature*
 361 *ecology & evolution* 2, 1556-1562. doi: 10.1038/s41559-018-0644-x [doi].

362 Braymer J. J. and Lill R. (2017). Iron-sulfur cluster biogenesis and trafficking in mitochondria. *The*
 363 *Journal of biological chemistry* 292, 12754-12763. doi: 10.1074/jbc.R117.787101 [doi].

364

365 Burki F, Kaplan M, Tikhonenkov DV, Zlatogursky V, Minh BQ, Radaykina LV, et al. Untangling the
 366 early diversification of eukaryotes: a phylogenomic study of the evolutionary origins of Centrohelida,
 367 Haptophyta and Cryptista. *Proc R Soc B Biol Sci.* 2016;283:20152802.

368 Couturier J., Touraine B., Briat J. F., Gaymard F. and Rouhier N. (2013). The iron-sulfur cluster
 369 assembly machineries in plants: current knowledge and open questions. *Frontiers in plant science* 4, 259.
 370 doi: 10.3389/fpls.2013.00259 [doi].

371 Danchin E. G. (2016). Lateral gene transfer in eukaryotes: tip of the iceberg or of the ice cube? *BMC*
 372 *biology* 14, 101-016-0330-x. doi: 10.1186/s12915-016-0330-x [doi].

373 Embley T. M. and Martin W. (2006). Eukaryotic evolution, changes and challenges. *Nature* 440, 623-
 374 630. doi: nature04546 [pii].

375 Embley T. M. and Williams T. A. (2015). Evolution: Steps on the road to eukaryotes. *Nature* 521, 169-
 376 170. doi: 10.1038/nature14522 [doi].

377 Eme L. and Ettema T. J. G. (2018). The eukaryotic ancestor shapes up. *Nature* 562, 352-353. doi:
 378 10.1038/d41586-018-06868-2 [doi].

379 Eme L., Spang A., Lombard J., Stairs C. W. and Ettema T. J. G. (2018). Archaea and the origin of
 380 eukaryotes. *Nature reviews.Microbiology* 16, 120. doi: 10.1038/nrmicro.2017.154 [doi].

381 Fontecave M., Choudens S. O., Py B. and Barras F. (2005). Mechanisms of iron-sulfur cluster assembly:
 382 the SUF machinery. *Journal of Biological Inorganic Chemistry : JBIC : a publication of the Society of*
 383 *Biological Inorganic Chemistry* 10, 713-721. doi: 10.1007/s00775-005-0025-1 [doi].

384 Freibert S. A., Goldberg A. V., Hacker C., Molik S., Dean P., Williams T. A., Nakjang S., Long S.,
 385 Sendra K., Bill E., Heinz E., Hirt R. P., Lucocq J. M., Embley T. M. and Lill R. (2017). Evolutionary
 386 conservation and in vitro reconstitution of microsporidian iron-sulfur cluster biosynthesis. *Nature*
 387 *communications* 8, 13932. doi: 10.1038/ncomms13932 [doi].

388 Gabaldon T. (2018). Relative timing of mitochondrial endosymbiosis and the "pre-mitochondrial
 389 symbioses" hypothesis. *IUBMB life* 70, 1188-1196. doi: 10.1002/iub.1950 [doi].

390 Gawryluk R. M. R. (2018). Evolutionary Biology: A New Home for the Powerhouse? *Current biology :*
 391 *CB* 28, R798-R800. doi: S0960-9822(18)30709-7 [pii].

392 Goldberg A. V., Molik S., Tsaousis A. D., Neumann K., Kuhnke G., Delbac F., Vivares C. P., Hirt R. P.,
 393 Lill R. and Embley T. M. (2008). Localization and functionality of microsporidian iron-sulfur cluster
 394 assembly proteins. *Nature* 452, 624-628. doi: 10.1038/nature06606 [doi].

395 Gray M. W., Burger G. and Lang B. F. (2001). The origin and early evolution of mitochondria. *Genome*
 396 *biology* 2, REVIEWS1018.

397 Gray M. W., Burger G. and Lang B. F. (1999). Mitochondrial evolution. *Science* 283, 1476-81.

398 Grosche C., Diehl A., Rensing S. A. and Maier U. G. (2018). Iron-Sulfur Cluster Biosynthesis in Algae
 399 with Complex Plastids. *Genome biology and evolution* 10, 2061-2071. doi: 10.1093/gbe/evy156 [doi].

400 Hirabayashi K., Yuda E., Tanaka N., Katayama S., Iwasaki K., Matsumoto T., Kurisu G., Outten F. W.,
 401 Fukuyama K., Takahashi Y. and Wada K. (2015). Functional Dynamics Revealed by the Structure of the
 402 SufBCD Complex, a Novel ATP-binding Cassette (ABC) Protein That Serves as a Scaffold for Iron-
 403 Sulfur Cluster Biogenesis. *The Journal of biological chemistry* 290, 29717-29731. doi:
 404 10.1074/jbc.M115.680934 [doi].

405 Hjort K., Goldberg A. V., Tsaousis A. D., Hirt R. P. and Embley T. M. (2010). Diversity and reductive
 406 evolution of mitochondria among microbial eukaryotes. *Philosophical transactions of the Royal Society*
 407 *of London.Series B, Biological sciences* 365, 713-727. doi: 10.1098/rstb.2009.0224 [doi].

408 Karnkowska A., Vacek V., Zubacova Z., Treitli S. C., Petrzalkova R., Eme L., Novak L., Zarsky V.,
 409 Barlow L. D., Herman E. K., Soukal P., Hroudova M., Dolezal P., Stairs C. W., Roger A. J., Elias M.,
 410 Dacks J. B., Vlcek C. and Hampl V. (2016). A Eukaryote without a Mitochondrial Organelle. *Current*
 411 *biology : CB* 26, 1274-1284. doi: 10.1016/j.cub.2016.03.053 [doi].

412 Koonin E. V. (2015). Origin of eukaryotes from within archaea, archaeal eukaryome and bursts of gene
 413 gain: eukaryogenesis just made easier? *Philosophical transactions of the Royal Society of London.Series*
 414 *B, Biological sciences* 370, 20140333. doi: 10.1098/rstb.2014.0333 [doi].

415 Lane N. and Martin W. F. (2016). Mitochondria, complexity, and evolutionary deficit spending.
 416 *Proceedings of the National Academy of Sciences of the United States of America* 113, E666. doi:
 417 10.1073/pnas.1522213113 [doi].

418 Lane N. and Martin W. F. (2015). Eukaryotes really are special, and mitochondria are why. *Proceedings*
 419 *of the National Academy of Sciences of the United States of America* 112, E4823. doi:
 420 10.1073/pnas.1509237112 [doi].

421 Le Corguille G., Pearson G., Valente M., Viegas C., Gschloessl B., Corre E., Bailly X., Peters A. F.,
 422 Jubin C., Vacherie B., Cock J. M. and Leblanc C. (2009). Plastid genomes of two brown algae,
 423 Ectocarpus siliculosus and Fucus vesiculosus: further insights on the evolution of red-algal derived
 424 plastids. *BMC evolutionary biology* 9, 253-2148-9-253. doi: 10.1186/1471-2148-9-253 [doi].

425 Leger M. M., Eme L., Hug L. A. and Roger A. J. (2016). Novel Hydrogenosomes in the Microaerophilic
 426 Jakobid Stygiella incarcerata. *Molecular biology and evolution* 33, 2318-2336. doi:
 427 10.1093/molbev/msw103 [doi].

428 Leger M. M., Eme L., Stairs C. W. and Roger A. J. (2018). Demystifying Eukaryote Lateral Gene
 429 Transfer (Response to Martin 2017 DOI: 10.1002/bies.201700115). *BioEssays : news and reviews in*
 430 *molecular, cellular and developmental biology* 40, e1700242. doi: 10.1002/bies.201700242 [doi].

431 Lill R. (2009). Function and biogenesis of iron-sulfur proteins. *Nature* 460, 831-838. doi:
 432 10.1038/nature08301 [doi].

433 Lill R., Diekert K., Kaut A., Lange H., Pelzer W., Prohl C. and Kispal G. (1999). The essential role of
 434 mitochondria in the biogenesis of cellular iron-sulfur proteins. *Biol Chem* 380, 1157-66.

435 Lill R., Dutkiewicz R., Freibert S. A., Heidenreich T., Mascarenhas J., Netz D. J., Paul V. D., Pierik A. J.,
 436 Richter N., Stumpfig M., Srinivasan V., Stehling O. and Muhlenhoff U. (2015). The role of mitochondria
 437 and the CIA machinery in the maturation of cytosolic and nuclear iron-sulfur proteins. *European journal*
 438 *of cell biology* 94, 280-291. doi: 10.1016/j.ejcb.2015.05.002 [doi].

439 Lill R., Hoffmann B., Molik S., Pierik A. J., Rietzschel N., Stehling O., Uzarska M. A., Webert H.,
 440 Wilbrecht C. and Muhlenhoff U. (2012). The role of mitochondria in cellular iron-sulfur protein
 441 biogenesis and iron metabolism. *Biochimica et biophysica acta* 1823, 1491-1508. doi:
 442 10.1016/j.bbamcr.2012.05.009 [doi].

443 Maralikova B., Ali V., Nakada-Tsukui K., Nozaki T., van der Giezen M., Henze K. and Tovar J. (2010).
 444 Bacterial-type oxygen detoxification and iron-sulfur cluster assembly in amoebal relict mitochondria.
 445 *Cellular microbiology* 12, 331-342. doi: 10.1111/j.1462-5822.2009.01397.x [doi].
 446 Margulis L., Chapman M., Guerrero R. and Hall J. (2006). The last eukaryotic common ancestor (LECA):
 447 Acquisition of cytoskeletal motility from aerotolerant spirochetes in the Proterozoic Eon. *Proc Natl Acad*
 448 *Sci USA* 103, 13080.
 449 Martijn J., Vosseberg J., Guy L., Offre P. and Ettema T. J. G. (2018). Deep mitochondrial origin outside
 450 the sampled alphaproteobacteria. *Nature* 557, 101-105. doi: 10.1038/s41586-018-0059-5 [doi].
 451 Martin W. F., Neukirchen S., Zimorski V., Gould S. B. and Sousa F. L. (2016). Energy for two: New
 452 archaeal lineages and the origin of mitochondria. *BioEssays : news and reviews in molecular, cellular*
 453 *and developmental biology* 38, 850-856. doi: 10.1002/bies.201600089 [doi].
 454 Mettert E. L., Outten F. W., Wanta B. and Kiley P. J. (2008). The impact of O(2) on the Fe-S cluster
 455 biogenesis requirements of Escherichia coli FNR. *Journal of Molecular Biology* 384, 798-811. doi:
 456 10.1016/j.jmb.2008.09.080 [doi].
 457 Miller C. N., Josse L. and Tsaousis A. D. (2018). Localization of Fe-S Biosynthesis Machinery in
 458 *Cryptosporidium parvum* Mitosome. *The Journal of eukaryotic microbiology* 65, 913-922. doi:
 459 10.1111/jeu.12663 [doi].
 460 Muller M. (1973). Peroxisomes and hydrogenosomes in protozoa. *The journal of histochemistry and*
 461 *cytochemistry : official journal of the Histochemistry Society* 21, 955-957. doi: 10.1177/21.11.955 [doi].
 462 Nyvltova E., Sutak R., Harant K., Sedinova M., Hrdy I., Paces J., Vlcek C. and Tachezy J. (2013). NIF-
 463 type iron-sulfur cluster assembly system is duplicated and distributed in the mitochondria and cytosol of
 464 *Mastigamoeba balamuthi*. *Proceedings of the National Academy of Sciences of the United States of*
 465 *America* 110, 7371-7376. doi: 10.1073/pnas.1219590110 [doi].

466 Outten F. W. (2015). Recent advances in the Suf Fe-S cluster biogenesis pathway: Beyond the
 467 Proteobacteria. *Biochimica et biophysica acta* 1853, 1464-1469. doi: 10.1016/j.bbamcr.2014.11.001
 468 [doi].

469 Paul V. D., Muhlenhoff U., Stumpfig M., Seebacher J., Kugler K. G., Renicke C., Taxis C., Gavin A. C.,
 470 Pierik A. J. and Lill R. (2015). The deca-GX3 proteins Yae1-Lto1 function as adaptors recruiting the
 471 ABC protein Rli1 for iron-sulfur cluster insertion. *eLife* 4, e08231. doi: 10.7554/eLife.08231 [doi].

472 Pena-Diaz P. and Lukes J. (2018). Fe-S cluster assembly in the supergroup Excavata. *Journal of*
 473 *Biological Inorganic Chemistry : JBIC : a publication of the Society of Biological Inorganic Chemistry*
 474 23, 521-541. doi: 10.1007/s00775-018-1556-6 [doi].

475 Pittis A. A. and Gabaldon T. (2016). Late acquisition of mitochondria by a host with chimaeric
 476 prokaryotic ancestry. *Nature* 531, 101-104. doi: 10.1038/nature16941 [doi].

477 Pyrih J., Pyrihova E., Kolisko M., Stojanovova D., Basu S., Harant K., Haindrich A. C., Dolezal P.,
 478 Lukes J., Roger A. and Tachezy J. (2016). Minimal cytosolic iron-sulfur cluster assembly machinery of
 479 *Giardia intestinalis* is partially associated with mitosomes. *Molecular microbiology* 102, 701-714. doi:
 480 10.1111/mmi.13487 [doi].

481 Rangachari K., Davis C. T., Eccleston J. F., Hirst E. M., Saldanha J. W., Strath M. and Wilson R. J.
 482 (2002). SufC hydrolyzes ATP and interacts with SufB from *Thermotoga maritima*. *FEBS letters* 514,
 483 225-228. doi: S0014579302023694 [pii].

484 Rouault T. A. and Maio N. (2017). Biogenesis and functions of mammalian iron-sulfur proteins in the
 485 regulation of iron homeostasis and pivotal metabolic pathways. *The Journal of biological chemistry* 292,
 486 12744-12753. doi: 10.1074/jbc.R117.789537 [doi].

487 Rudolf J., Makrantonis V., Ingledew W. J., Stark M. J. and White M. F. (2006). The DNA repair helicases
 488 XPD and FancJ have essential iron-sulfur domains. *Molecular cell* 23, 801-808. doi: S1097-
 489 2765(06)00516-8 [pii].

490 Santos H. J., Makiuchi T. and Nozaki T. (2018). Reinventing an Organelle: The Reduced Mitochondrion
 491 in Parasitic Protists. *Trends in parasitology* 34, 1038-1055. doi: S1471-4922(18)30173-9 [pii].
 492 Sousa F. L., Neukirchen S., Allen J. F., Lane N. and Martin W. F. (2016). Lokiarchaeon is hydrogen
 493 dependent. *Nature microbiology* 1, 16034. doi: 10.1038/nmicrobiol.2016.34 [doi].
 494 Spang A., Caceres E. F. and Ettema T. J. G. (2017). Genomic exploration of the diversity, ecology, and
 495 evolution of the archaeal domain of life. *Science (New York, N.Y.)* 357, 10.1126/science.aaf3883. doi:
 496 eaaf3883 [pii].
 497 Spang A., Eme L., Saw J. H., Caceres E. F., Zaremba-Niedzwiedzka K., Lombard J., Guy L. and Ettema
 498 T. J. G. (2018). Asgard archaea are the closest prokaryotic relatives of eukaryotes. *PLoS genetics* 14,
 499 e1007080. doi: 10.1371/journal.pgen.1007080 [doi].
 500 Spang A. and Ettema T. J. G. (2017). Archaeal evolution: The methanogenic roots of Archaea. *Nature*
 501 *microbiology* 2, 17109. doi: 10.1038/nmicrobiol.2017.109 [doi].
 502 Spang A., Saw J. H., Jorgensen S. L., Zaremba-Niedzwiedzka K., Martijn J., Lind A. E., van Eijk R.,
 503 Schleper C., Guy L. and Ettema T. J. G. (2015). Complex archaea that bridge the gap between
 504 prokaryotes and eukaryotes. *Nature* 521, 173-179. doi: 10.1038/nature14447 [doi].
 505 Stairs C. W., Eme L., Brown M. W., Mutsaers C., Susko E., Dellaire G., Soanes D. M., van der Giezen
 506 M. and Roger A. J. (2014). A SUF Fe-S cluster biogenesis system in the mitochondrion-related organelles
 507 of the anaerobic protist *Pygusua*. *Current biology : CB* 24, 1176-1186. doi: 10.1016/j.cub.2014.04.033
 508 [doi].
 509 Stehling O., Mascarenhas J., Vashisht A. A., Sheftel A. D., Niggemeyer B., Rosser R., Pierik A. J.,
 510 Wohlschlegel J. A. and Lill R. (2013). Human CIA2A-FAM96A and CIA2B-FAM96B integrate iron
 511 homeostasis and maturation of different subsets of cytosolic-nuclear iron-sulfur proteins. *Cell metabolism*
 512 18, 187-198. doi: 10.1016/j.cmet.2013.06.015 [doi].

513 Stehling O., Vashisht A. A., Mascarenhas J., Jonsson Z. O., Sharma T., Netz D. J., Pierik A. J.,
514 Wohlschlegel J. A. and Lill R. (2012). MMS19 assembles iron-sulfur proteins required for DNA
515 metabolism and genomic integrity. *Science (New York, N.Y.)* 337, 195-199. doi: 10.1126/science.1219723
516 [doi].

517 Stehling O., Wilbrecht C. and Lill R. (2014). Mitochondrial iron-sulfur protein biogenesis and human
518 disease. *Biochimie* 100, 61-77. doi: 10.1016/j.biochi.2014.01.010 [doi].

519 Tokumoto U., Kitamura S., Fukuyama K. and Takahashi Y. (2004). Interchangeability and distinct
520 properties of bacterial Fe-S cluster assembly systems: functional replacement of the isc and suf operons in
521 *Escherichia coli* with the nifSU-like operon from *Helicobacter pylori*. *Journal of Biochemistry* 136, 199-
522 209. doi: 10.1093/jb/136.2.199 [pii].

523 Tonini M. L., Pena-Diaz P., Haindrich A. C., Basu S., Kriegova E., Pierik A. J., Lill R., MacNeill S. A.,
524 Smith T. K. and Lukes J. (2018). Branched late-steps of the cytosolic iron-sulfur cluster assembly
525 machinery of *Trypanosoma brucei*. *PLoS pathogens* 14, e1007326. doi: 10.1371/journal.ppat.1007326
526 [doi].

527 Tovar J., Fischer A. and Clark C. G. (1999). The mitosome, a novel organelle related to mitochondria in
528 the amitochondrial parasite *Entamoeba histolytica*. *Mol Microbiol* 32, 1013-21.

529 Tovar J., Leon-Avila G., Sanchez L. B., Sutak R., Tachezy J., van der Giezen M., Hernandez M., Muller
530 M. and Lucocq J. M. (2003). Mitochondrial remnant organelles of *Giardia* function in iron-sulfur protein
531 maturation. *Nature* 426, 172-6.

532 Treitli S. C., Kotyk M., Yubuki N., Jirounkova E., Vlasakova J., Smejkalova P., Sipek P., Cepicka I. and
533 Hampl V. (2018). Molecular and Morphological Diversity of the Oxymonad Genera *Monocercomonoides*
534 and *Blattamonas* gen. nov. *Protist* 169, 744-783. doi: 10.1016/j.protist.2018.05.003 [pii].

535 Tsaousis A. D., Gentekaki E., Eme L., Gaston D. and Roger A. J. (2014). Evolution of the cytosolic iron-
536 sulfur cluster assembly machinery in *Blastocystis* species and other microbial eukaryotes. *Eukaryotic cell*
537 13, 143-153. doi: 10.1128/EC.00158-13 [doi].

538 Tsaousis A. D., Kunji E. R., Goldberg A. V., Lucocq J. M., Hirt R. P. and Embley T. M. (2008). A novel
539 route for ATP acquisition by the remnant mitochondria of *Encephalitozoon cuniculi*. *Nature* 453, 553-
540 556. doi: 10.1038/nature06903 [doi].

541 Tsaousis A. D., Ollagnier de Choudens S., Gentekaki E., Long S., Gaston D., Stechmann A., Vinella D.,
542 Py B., Fontecave M., Barras F., Lukes J. and Roger A. J. (2012). Evolution of Fe/S cluster biogenesis in
543 the anaerobic parasite *Blastocystis*. *Proceedings of the National Academy of Sciences of the United States*
544 *of America* 109, 10426-10431. doi: 10.1073/pnas.1116067109 [doi].

545 Vacek V., Novak L. V. F., Treitli S. C., Taborsky P., Cepicka I., Kolisko M., Keeling P. J. and Hampl V.
546 (2018). Fe-S Cluster Assembly in Oxymonads and Related Protists. *Molecular biology and evolution* 35,
547 2712-2718. doi: 10.1093/molbev/msy168 [doi].

548 van der Giezen M., Cox S. and Tovar J. (2004). The iron-sulfur cluster assembly genes *iscS* and *iscU* of
549 *Entamoeba histolytica* were acquired by horizontal gene transfer. *BMC evolutionary biology* 4, 7-2148-4-
550 7. doi: 10.1186/1471-2148-4-7 [doi].

551 Williams B. A., Hirt R. P., Lucocq J. M. and Embley T. M. (2002). A mitochondrial remnant in the
552 microsporidian *Trachipleistophora hominis*. *Nature* 418, 865-9.

553 Wollers S., Layer G., Garcia-Serres R., Signor L., Clemancey M., Latour J. M., Fontecave M. and
554 Ollagnier de Choudens S. (2010). Iron-sulfur (Fe-S) cluster assembly: the SufBCD complex is a new type
555 of Fe-S scaffold with a flavin redox cofactor. *The Journal of biological chemistry* 285, 23331-23341. doi:
556 10.1074/jbc.M110.127449 [doi].

557 Zaremba-Niedzwiedzka K., Caceres E. F., Saw J. H., Backstrom D., Juzokaite L., Vancaester E., Seitz K.
558 W., Anantharaman K., Starnawski P., Kjeldsen K. U., Stott M. B., Nunoura T., Banfield J. F., Schramm

A., Baker B. J., Spang A. and Ettema T. J. (2017). Asgard archaea illuminate the origin of eukaryotic cellular complexity. *Nature* 541, 353-358. doi: 10.1038/nature21031 [doi].

Figure legends:

Figure 1: Cartoon model of the mitochondrial Fe/S protein assembly process.

Figure was produced based on Braymer and Lill (2017). A cascade of ISC proteins is required for the de novo synthesis of [2Fe-2S] and [4Fe-4S] clusters and their proper trafficking to target apoproteins in mitochondria. Initially, a [2Fe-2S] cluster is synthesized by the early ISC machinery, composed of the Isu1 scaffold protein requiring sulfide from the cysteine desulfurase complex Nfs1-Isd11-Acp1, electrons from the transfer chain NADPH-Arh1 and the ferredoxin Yah1, and the regulator and/or iron donor Yfh1. The Isu1-bound [2Fe-2S] cluster is then delivered to the monothiol glutaredoxin Grx5, a reaction accomplished by the Hsp70 chaperone Ssq1 with the help of the J-type co-chaperone Jac1. This reaction is dependent on ATP hydrolysis by Ssq1. The exchange factor Mge1 facilitates the exchange of ADP for ATP. The resulting bridging [2Fe-2S] cluster on a Grx5 dimer is inserted directly into [2Fe-2S] recipient apoproteins or trafficked to the late ISC machinery for [4Fe-4S] cluster biogenesis. The early ISC machinery, including the chaperones and Grx5, is also responsible for generating the component X-S for transport of sulfur out of the mitochondria to the CIA machinery for cytosolic-nuclear Fe/S protein biogenesis. The late ISC machinery consists of the yet structurally and functionally uncharacterized Isa1-Isa2-Iba57 complex and is needed for the generation of [4Fe-4S] clusters. Trafficking and insertion of the [4Fe-4S] clusters into target Fe/S proteins are facilitated by specific ISC targeting factors, such as Nfu1, the complex I-specific Ind1, and the Bol proteins. Dashed arrows indicate steps that remain poorly elucidated on the biochemical level.

Figure 2: Cartoon demonstrating the current model, based on Braymer and Lill (2017), for the mechanism of yeast cytosolic-nuclear Fe-S protein biogenesis (a) and a hypothetical model for the *Blastocystis* (b) and the amitochondriate *Monocercomonoides* (c).

Assembly of extra-mitochondrial Fe-S proteins is catalyzed by the cytosolic iron–sulfur protein assembly (CIA) machinery in an ISC-dependent manner. Several CIA protein complexes support different stages of the process. Initially, a bridging [4Fe–4S] cluster is assembled on the Cfd1–Nbp35 scaffold complex, but the bridging cluster binds only transiently. Nbp35 contains another stably bound [4Fe–4S] cluster at its N-terminus. Cluster assembly on Cfd1–Nbp35 depends on the molecule X–S from the mitochondrial ISC machinery. Further, the electron transfer chain from NADPH via the diflavin reductase Tah18 and the Fe–S protein Dre2 is needed. In a second step, the transiently bound [4Fe–4S] cluster of Cfd1–Nbp35 is transferred to and inserted into apoproteins by the Fe–S protein Nar1, and the CIA targeting complex consisting of Cia1, Cia2 and Mms19. Maturation of the essential Fe–S protein Rli1 additionally depends on the function of the two specific adaptor proteins Yae1 and Lto1. The Yae1–Lto1 complex uses a unique binding cascade to recruit Rli1 to the CIA targeting complex for Fe–S cluster insertion.

Figure 3: The distribution of the SUF system amongst microbes and scenarios on the evolution of the SUF machinery in eukaryotes

a. The distribution of the SUF system amongst microbial genomes [based on Tokumoto et al. (2004)]. Since the sufBC-like genes are found in all species encoding this system, it has been speculated that these genes were components of the primitive system, which was further evolved through the recruitment of other components such as SufA, SufE and SufS (e.g. *E. coli* Suf system). The fused genes found in *Blastocystis*, *Pygusua* and *Stygiella* genomes/transcriptoms corresponding to the SufCB operon in methanomicrobiales. The SufCB operon encodes two out of the six proteins of the SUF system (e.g. *E. coli* or plastid bearing organisms) and is part of the Suf system found in extremophiles. **b.** The eukaryotic tree of life demonstrating the distribution of the various Fe/S cluster biosynthetic pathways in eukaryotes, highlighting (purple color) the unique distribution of the SUF system across eukaryotes. Relationships between eukaryotes are based on recent concatenated phylogenetic results (Burki et al. 2016). **c.** This scenario suggests that the common ancestor of *Blastocystis* has acquired the fused gene from a methanoarchaeon, while *Pygusua* and *Stygiella* independently acquiring the SufCB fused gene from an organism from the same group of methanomicrobiales as well. **d.** In this scenario, the last common ancestor of *Blastocystis* acquired the SufCB fused gene from an organism from the group of

614 methanomicrobiales which was laterally gene transferred to *Pygsuia* and *Stygiella*. **e.** In this scenario
615 *Stygiella* acquired the SufCB fused gene from an organism from the group of methanomicrobiales which
616 was laterally gene transferred to *Pygsuia* and the last common ancestor of *Blastocystis*. **f.** In this scenario
617 *Pygsuia* acquired the SufCB fused gene from an organism from the group of methanomicrobiales which
618 was laterally gene transferred to *Stygiella* and the last common ancestor of *Blastocystis*. **g.** In this
619 scenario, the methanoarchaeal SufCB was either present in last eukaryotic common ancestor or was
620 acquired later before the split of the various eukaryotic lineages.

In review

Figure 1.JPEG




Figure 2.JPEG




Figure 3.JPEG

